



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07D 409/04, 405/04, 333/38, 307/68, C07C 327/58	A1	(11) International Publication Number: WO 00/24738 (43) International Publication Date: 4 May 2000 (04.05.00)
<p>(21) International Application Number: PCT/US99/24860</p> <p>(22) International Filing Date: 22 October 1999 (22.10.99)</p> <p>(30) Priority Data: 60/105,314 23 October 1998 (23.10.98) US</p> <p>(71) Applicant: DOW AGROSCIENCES LLC [US/US]; 9330 Zionsville Road, Indianapolis, IN 46268 (US).</p> <p>(72) Inventors: ASH, Mary, L.; 762 Franklin Trace, Zionsville, IN 46077 (US). ZETTLER, Mark, W.; 11142 St. Charles Place, Carmel, IN 46033 (US). PEARSON, Norman, R.; 9622 Turnberry Court, Carmel, IN 46032 (US). ROMER, Duane, R.; 4025 Lowell Court, Midland, MI 48642 (US). HULL, John, W., Jr.; 3104 Saint Mary's Drive, Midland, MI 48640 (US). PODHOREZ, David, E.; 5604 Berry Court, Midland, MI 48640 (US).</p> <p>(74) Agent: STUART, Donald, R.; Dow Agrosiences LLC, 9330 Zionsville Road, Indianapolis, IN 46268 (US).</p>		<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: PROCESS FOR PREPARING 3-(SUBSTITUTED PHENYL)-5-THIENYL OR FURYL-1,2,4-TRIAZOLES AND NOVEL INTERMEDIATES UTILIZED THEREIN</p> <p>(57) Abstract</p> <p>New synthetic procedures for preparing insecticidal 3-(substituted phenyl)-5-(thienyl or furyl)-1,2,4-triazoles utilizing thioimide intermediate.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.


AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

PROCESS FOR PREPARING 3-(SUBSTITUTED PHENYL)-5-THIENYL OR
FURYL)-1,2,4-TRIAZOLES AND NOVEL INTERMEDIATES UTILIZED
THEREIN

5 This invention provides new synthetic procedures and intermediates for preparing insecticidal 3-(substituted phenyl)-5-thienyl or furyl)-1,2,4-triazoles.

WO 98/47894 discloses a series of highly active new insecticides. Although that patent application discloses
10 suitable laboratory methods for preparation of the compounds, a need exists for a commercially applicable manufacturing process.

The present invention provides a process for preparing compounds of the formula (1)




15

wherein

Ar is substituted phenyl;

Y is O or S;

R^3 is selected from H, halo, lower alkyl, (C₇-C₂₁)
20 straight or branched chain alkyl, hydroxy, lower alkoxy, haloalkyl, haloalkoxy, alkoxyalkyl, alkoxyalkoxy, lower alkenyl, lower alkynyl, haloalkenyl, CN, NO₂, COR⁶, CO₂R⁶, CON(R⁶)₂, (C₃-C₆) cycloalkyl, S(O)_mR⁶, -OSO₂R⁶, SCN, -
(CH₂)_nR⁶, -CH=CHR⁶, -C≡CR⁶, -(CH₂)_qOR⁶, -(CH₂)_qSR⁶, -
25 (CH₂)_qNR⁶R⁶, -O(CH₂)_qR⁶, -S(CH₂)_qR⁶, -NR⁶(CH₂)_qR⁶,



$\text{—NR}^6\text{C(=O)N(R}^6\text{)}$, $\text{—C(=NOR}^6\text{)R}^6$, —OC(=O)R^6 , or —SC(=O)R^6 , pyridyl, substituted pyridyl, isoxazolyl, substituted isoxazolyl, naphthyl, substituted naphthyl, phenyl, substituted phenyl, thienyl, substituted thienyl, pyrimidyl, substituted pyrimidyl, pyrazolyl, or substituted pyrazolyl;

R^4 and R^5 are independently H, halo, lower alkyl, lower alkoxy, haloalkyl, haloalkoxy, CN, CO_2R^6 , $\text{CON(R}^6\text{)}_2$, or S(O)_m alkyl, or

if R^4 and R^5 are attached to adjacent carbon atoms, they may join to form a 5 or 6 member saturated or unsaturated carbocyclic ring which may be substituted by 1 or 2 halo, lower alkyl, lower alkoxy or haloalkyl groups;

R^6 is H, lower alkyl, haloalkyl, lower alkenyl, lower alkynyl, phenyl, or substituted phenyl;

m is 0, 1, or 2;


n is 1 or 2;

p is an integer from 2 to 6; and


q is 0 or 1;

which comprises the steps of:


(a) reacting a compound of formula (2)



wherein Ar is as defined in formula (1) and R^7 is lower alkyl, or an acid addition salt thereof, with an acid chloride of the formula (3)



wherein Y, R³, R⁴, and R⁵ are as defined in formula (1),
 in an inert organic solvent the presence of an organic or
 inorganic base to produce the adduct-intermediate of
 5 formula (4)




wherein Ar, Y, R³, R⁴, and R⁵ are as defined in formula
 (1) and R⁷ is lower alkyl; and

(b) optionally isolating said adduct intermediate,
 10 and

(c) reacting said adduct-intermediate with methyl
 hydrazine to produce the compound of formula (1).

Adduct-intermediates of formula (4) are novel
 compounds, and they are an important aspect of the
 15 invention.

Adduct-intermediates that are preferred based on
 desirable properties of the final products they may be
 used to produce are those of formula



20

wherein

R¹ and R² are independently F or Cl;


R³, R⁴, and R⁵ are independently H, CH₃, Cl or Br;
and

R⁷ is lower alkyl.

Most preferred adduct intermediates are those of
5 formula (4a) wherein R¹ is Cl, R² is F, R⁷ is CH₃, and:

- (a) R³, R⁴, and R⁵ are each Cl;
- (b) R³ and R⁴ are each Br and R⁵ is H; or
- (c) R³ is CH₃, R⁴ is Cl or Br, and R⁵ is H.

In another of its aspects, the invention provides
10 the methyl sulfate salt of a compound of the formula



Throughout this document, all temperatures are given
in degrees Celsius, and all percentages are weight
percentages unless otherwise stated.

15 The term "lower alkyl" refers to (C₁-C₆) straight
hydrocarbon chains and (C₃-C₆) branched and cyclic
hydrocarbon groups.

The terms "lower alkenyl" and "lower alkynyl" refer
to (C₂-C₆) straight hydrocarbon chains and (C₃-C₆)
20 branched hydrocarbon groups containing at least one
double or triple bond, respectively.

The term "lower alkoxy" refers to -O-lower alkyl.

The terms "halomethyl" and "haloalkyl" refer to
methoxy and lower alkyl groups substituted with one or
25 more halo atoms.

The terms "halomethoxy" and "haloalkoxy" refer to methyl and lower alkoxy groups substituted with one or more halo atoms.

5 The term "alkoxyalkyl" refers to a lower alkyl group substituted with a lower alkoxy group.

The terms "substituted naphthyl", "substituted thienyl," "substituted pyrimidyl," "substituted pyrazolyl," "substituted pyridyl," and "substituted isoxaxolyl" refer to the ring system substituted with one
10 or more groups independently selected from halo, halo (C₁-C₄) alkyl, CN, NO₂, (C₁-C₄) alkyl, (C₃-C₄) branched alkyl, phenyl, (C₁-C₄) alkoxy, or halo (C₁-C₄) alkoxy.

The term "substituted phenyl" refers to a phenyl group substituted with one or more groups independently
15 selected from halo, (C₁-C₁₀) alkyl, branched (C₃-C₆) alkyl, halo (C₁-C₇) alkyl, hydroxy (C₁-C₇) alkyl, (C₁-C₇) alkoxy, halo (C₁-C₇) alkoxy, phenoxy, phenyl, NO₂, OH, CN, (C₁-C₄) alkanoyl, benzoyl, (C₁-C₄) alkanoyloxy, (C₁-C₄) alkoxy carbonyl, phenoxy carbonyl, or benzoyloxy.

20 The term "substituted benzenesulfonyl" refers to p-chlorobenzenesulfonyl and p-toluenesulfonyl.

Unless otherwise indicated, when it is stated that a group may be substituted with one or more substituents selected from an identified class, it is intended that
25 the substituents may be independently selected from the class.

Alkyl benzthioimidates of formula (2) are known in the literature. They are preferably used in the present invention as an acid addition salt. For example,
30 tetrafluoroboric acid, hydrogen chloride, hydrogen bromide, hydrogen iodide, or the like, may be used. The methyl sulfate salt is preferred.


Thioimidates are readily available through alkylation of the corresponding thioamides which are themselves commercially available or can be made from the amide (Phosphorus Sulfur (1985), 25(3), 297-305) or nitrile (Chem.-Ztg. (1980), 104(12), 365-7; J. Chem. Soc. (1952), 742; Can. J. Chem. (1985), 63, 3075).

Reaction of the acid chloride (3) and the imidate (2) to give the adduct (4) can generally be accomplished in any inert solvent with any organic or inorganic base.

Reaction of compounds of formula (4) with methyl hydrazine gives the desired triazoles, generally in good yield and with a high degree of regiospecificity. An example of a previous application of a similar process is given in Synthesis, 483 (1983). In regard to regiospecificity, the undesired possible coproducts are the 5-(substituted phenyl)-3-thienyl or furyl isomers of the desired 3-(substituted phenyl)-5-thienyl or furyl-1,2,4-triazoles.

Compounds of formula (1) can also be made starting with an alkyl benzimidate, as illustrated in the following comparative example. It has been found, however, that the process of the invention, utilizing alkyl benzthioimidates, provides significantly better yields.

25 Comparative Example 1



Briefly, the reaction of ethyl 2-fluoro-6-chlorobenzimidate (5) with 3,4,5-trichloro-2-thienoyl chloride (3a) was performed in toluene using

triethylamine as an acid scavenger. The presumed intermediate (6) which was not isolated, then reacted with methyl hydrazine 3 hr at reflux to give 1-methyl-3-(2-chloro-6-fluoro)phenyl-5-(3,4,5-trichloro-2-thienyl)-1H-1,2,4-triazole (1a) in 35% yield. The following paragraphs describe each step of the synthesis in greater detail.

A. 2-chloro-6-fluorobenzamide

The procedure of R. Balicki and L. Kaczmarek, Syn. Comm., **23**, 3149 (1993) was used. Into a 1 L round bottom flask equipped with a magnetic stirrer was added acetone (60 mL), water (60 mL), 2-chloro-6-fluorobenzonitrile (14.5 g, 93.5 mmol), urea (22.5 g, 374 mmol), 30% hydrogen peroxide (42.4 g, 374 mmol), and potassium carbonate (1.3 g, 9.0 mmol). The resulting slurry was stirred at room temperature overnight. Additional portions of urea (11.2 g, 30% hydrogen peroxide (41 g), acetone (30 mL) and potassium carbonate (0.63 g) were added and the mixture was allowed to stir for an additional 2 hr. Additional portions of urea (11.2 g, 30% hydrogen peroxide (41 g), acetone (30 mL) and potassium carbonate (0.63 g) were again added and the mixture was allowed to stir overnight. The acetone was removed in vacuo, the residue extracted with methylene chloride, the organics were combined, washed with brine, dried (MgSO₄) and the solvent removed in vacuo to give 15.8 g (97.5% yield) of 2-chloro-6-fluorobenzamide (97% pure by GC): mp 140-141 °C; ¹H NMR (CDCl₃) δ 7.2-7.4 (m, 2H), 7.0 (m, 1H), 6.3 (s, b, 1H), 5.9 (s, b, 1H).

B. Ethyl 2-chloro-6-fluorobenzimidate (5)

Into a 1 L round bottom flask equipped with a mechanical stirrer, thermometer, condenser, dropping funnel, under an atmosphere of nitrogen was added methylene chloride (240 mL) and 2-chloro-6-fluorobenzamide (8.65 g, 50 mmol). The


temperature of the mixture was cooled to approximately 0 °C and a solution of triethyloxonium tetrafluoroborate (55 mL of 1M soln) was added dropwise at a rate such that the temperature did not rise over 0 °C. The resulting mixture was allowed to warm to stir cold for 1 hr , the cooling bath was removed, the mixture was allowed to warm to RT, and was stirred at that temperature overnight. Due to incomplete conversion the mixture was cooled to °5 C and dropwise was added and additional 11 mL of 1M soln of triethyloxonium tetrafluoroborate. The cooling bath was removed and the mixture allowed to stir for an additional 24 hr. The mixture was cooled to 0 °C and an additional portion of triethyloxonium tetrafluoroborate (10 mL) was added dropwise, the bath removed and the mixture stirred at RT for 19 hr. The temperature was lowered to 0 °C and a solution of potassium carbonate (50%) was added dropwise. After stirring for 15 min the layers were separated, the aqueous layer washed with methylene chloride, the organics combined, dried (Na₂SO₄), and the solvent removed in vacuo to give an oil: ethyl 2-chloro-6-fluorobenzimidate (94%), starting material (5%). This material was used without further purification.

C. Preparation of 1-methyl-3-(2-chloro-6-fluoro)phenyl-5-(3,4,5-trichloro-2-thienyl)-1H-1,2,4-triazole(1a) from ethyl 2-chloro-6-fluorobenzimidate (5)

Into 50 mL one necked round bottom flask equipped with a magnetic stirrer and condenser under an atmosphere of nitrogen was added 3,4,5-trichloro-2-thienyl carboxylic acid (0.69 g, 3 mmol), EDC (20 mL), thionyl chloride (0.53 g, 0.32 mL, 4.5 mmol) and several drops of DMF. The mixture was allowed to reflux for 3 hr and the solvent was removed in vacuo to a constant weight. To a three necked flask equipped with a condenser and containing ethyl 2-chloro-6-fluorobenzimidate (94% pure, 0.60 g, 3.0 mmol), and triethylamine (0.10 g, 0.14 mL,

6.0 mmol), and toluene (10 mL) was added dropwise 3,4,5-trichloro-2-thienoyl chloride dissolved in toluene (10 mL). The temperature of the mixture was brought to the point of reflux and maintained at that temperature for 3 hr. To the refluxing mixture was added methyl hydrazine (0.86 g, 1.0 mL, 9 mmol) dissolved in toluene was added dropwise. The mixture was allowed to reflux for 1 hr. The toluene was removed in vacuo and the residue was purified via column chromatography to give 1-methyl-3-(2-chloro-6-fluoro)phenyl-5-(3,4,5-trichloro-2-thienyl)-1H-1,2,4-triazole (**1a**): 0.42 g (35% yield), GC[Mez80] at 19.1 min, mass 395/397/399.

Example 1



Briefly, the reaction of methyl 2-fluoro-6-chlorobenzthioimidate hydroiodide (**2a**) with 3,4,5-trichloro-2-thienoyl chloride (**3a**) was performed in toluene using triethylamine as an acid scavenger. The presumed intermediate (**4a**) which was not isolated, then reacted with methyl hydrazine for 3 hr at reflux to give 1-methyl-3-(2-chloro-6-fluoro)phenyl-5-(3,4,5-trichloro-2-thienyl)-1H-1,2,4-triazole (**1a**) in 78% yield.

The following paragraphs describe each step of the synthesis in greater detail.

A. Preparation of 2-chloro-6-fluorobenzthioamide

Into a 250-mL three necked round bottom flask equipped with a mechanical stirrer, dry ice condenser, dropping funnel, and outlet to a trap filled with bleach was added pyridine (45 mL), 2-chloro-6-fluorobenzonitrile (15.5 g,

0.1 mol), triethylamine (13.4 g, 18.6 mL, 0.133 mol), and sodium sulfide hydrate (36.0 g, 0.15 mol). The temperature of the resulting solution was lowered to approximately 5 °C using an ice bath. To the stirred
5 slurry was added dropwise concentrated hydrochloric acid (25.8 g, 0.266 mol). The rate of addition of the hydrochloric acid was such that the temperature of the reaction mixture did not exceed 25 °C with an addition time of 20 min. The cooling bath was removed and the
10 slurry was allowed to warm to RT and to stir over night. An additional 40 grams of sodium sulfide and 50 mL of hydrochloric acid was added to the mixture and allowed to stir over night. An additional 40 grams of sodium sulfide and 50 mL of hydrochloric acid was added and the
15 mixture allowed to stir over the weekend. The mixture was poured into water (2 L) and was extracted with ether. The ether layer was washed with dilute sulfuric acid, water, brine, dried (MgSO₄), and the solvent removed in vacuo to give 16.1 g of crude product which was
20 recrystallized from ethyl acetate to give 2-chloro-6-fluoro-benzthioamide as light yellow crystals: mp 156-158 °C.

B. Preparation of methyl 2-chloro-6-fluorobenzthioimide hydroiodide (2a)

25 Into a 50 mL three necked round bottom flask equipped with a magnetic stirrer was added 2-chloro-6-fluorobenzthioamide (3.78 g, 20 mmol) and acetone (20 mL). Dropwise was added iodomethane (2.84 g, 1.24 mL, 20 mmol) and the mixture was allowed to stir at room
30 temperature overnight with monitoring via HPCL (60/40 acetonitrile/water). The solids were removed via filtration and washed with small portions of acetone to give methyl 2-chloro-6-fluorobenzthioimide hydroiodide as a light yellow solid: 3.1 g (46.9% yield): ¹H NMR
35 (DMSO-d₆) δ 7.7 (m, 1H), 7.5 (m, 2H), 2.7 (s, 3H).

C. Preparation of 1-methyl-3-(2-chloro-6-fluoro)phenyl-5-(3,4,5-trichloro-2-thienyl)-1H-1,2,4-triazole (1a) from 2-chloro-6-fluorobenzthioimide hydroiodide (2a)


Into 50 mL one necked round bottom flask equipped with a
5 magnetic stirrer and condenser under an atmosphere of
nitrogen was added 3,4,5-trichloro-2-thienyl carboxylic
acid (0.69 g, 3 mmol), EDC (20 mL), thionyl chloride
(0.53 g, 0.32 mL, 4.5 mmol) and several drops of DMF.
The mixture was allowed to reflux for 3 hr and the
10 solvent was removed in vacuo to a constant weight. To a
three necked flask equipped with a condenser and
containing methyl 2-chloro-6-fluorobenzthioimide
hydroiodide (0.99 g, 3.0 mmol), and triethylamine (0.20
g, 0.28 mL, 12 mmol), and toluene (20 mL) was added
15 dropwise 3,4,5-trichloro-2-thienoyl chloride dissolved in
toluene (10 mL). The mixture was brought to the point of
reflux and maintained at that temperature for 3 hr. To
the refluxing mixture was added methyl hydrazine (0.86 g,
1.0 mL, 18 mmol) dissolved in toluene was added dropwise.
20 The mixture was allowed to reflux for 1 hr, the toluene
was removed in vacuo, and the residue chromatographed on
silica gel to give 0.87 g (73% yield), GC[Mez80] at 19.1
min, mass 395/397/399.

Thus, the process of the invention provided a
25 significantly better yield (73% versus 35%) than the
process using an alkyl benzimidate in place of the alkyl
benzthioimide required by the invention.

The preferred process for preparing the methyl
2-chloro-6-fluorobenzthioimide produces the methyl
30 sulfate salt.

Example 2

Preparation of methyl 2-chloro-6-fluorobenzthioimide
using dimethylsulfate



Into a 25-mL three-necked round bottom flask equipped with a magnetic stirrer was added 2-chloro-6-fluorobenzthioamide (1.89 g, 10.0 mmol) and toluene (10 mL). Dimethylsulfate was added dropwise and the mixture was allowed to stir at reflux using a Dean Stark trap. The solids were removed via filtration and washed with small portions of toluene to give methyl 2-chloro-6-fluorobenzthioimidate as a light yellow solid: 2.7 g (87% yield): ^1H NMR (DMSO- d_6) δ 7.7 (m, 1H), 7.5 (m, 2H), 3.3 (s, 3H), 2.7 (s, 3H); 98% pure by HPLC analysis.

Comparing the results of Example 2 with those of Example 1C, it is clear that Example 2 is a significant improvement. Example 2 is higher yielding (87% versus 46%), yields a purer product (98% versus 96%) and uses a reagent, dimethyl sulfate, that is much cheaper both in terms of initial cost and disposal issues.

The following example illustrates the complete process incorporating use of the methyl sulfate salt and isolation of the adduct intermediate.

Example 3

A. 2-Chloro-6-Fluorobenzthioamide

2-Chloro-6-fluorobenzonitrile, (99.1%, 62.2 g, 0.40 mol) was weighed into a 1-L three-necked roundbottom flask equipped with a condenser and an overhead electric stirrer, along with Et_3N (78 mL, 56.6 g, 0.56 mol) and 180 mL (176.04 g, 2.23 mol) of pyridine. The reactor was purged with a slow stream of N_2 and vented to a 13% bleach solution. The stirring solution was cooled to -19°C in a CCl_4 /dry ice bath, and H_2S gas (33.6 g, 0.99 mol) was

sparged below the liquid surface at a rate of 0.4 g/min over a period of 82 min. During the gas addition, the solution temperature rose to -11°C. The yellow-green solution was allowed to gradually warm to 25°C and stir
5 overnight with a slow N₂ purge of the reactor head space into bleach solution. The solution was poured into 1.6 L of ice water, stirred, and the resulting white crystals were collected on a buchner funnel and rinsed with additional water. After 2 h of air drying, the moist
10 filter cake was vacuum oven dried for 5 h at 65°C to give 54.5 g of 2-Chloro-6-Fluorobenzthioamide (72% wt. % yield), mp 155-160°C, having a GC area % purity of 98.8 % and containing 1.2 % 2-Chloro-6-fluorobenzonitrile.

15 B. Methyl 2-chloro-6-fluorobenzthioimide methyl sulfate salt


2-Chloro-6-fluorobenzthioamide (21.9 g, 0.12 mol) and dimethyl sulfate (12 mL, 16.0 g, 0.13 mol) were stirred together in 100 mL of toluene and refluxed for 2 h. The slurry was cooled to 25°C and the white crystals
20 collected on a glass buchner funnel and rinsed with a small quantity of acetone. After air suction drying for 1 h, 35.5 g of methyl 2-chloro-6-fluorobenzthioimide methylsulfate salt was obtained, mp 147-151°C, representing a nominal wt. % yield of 96%.

25 A sample of this salt was recrystallized from cold isopropanol followed by vacuum oven drying at 75°C to give white crystals, mp 152-157°C. Elemental analysis results are summarized in the table below.

	<u>Theoretical</u>	<u>Found</u>
C	34.23%	34.47%
H	3.51%	3.61%
N	4.44%	4.45%

Cl	11.23%	11.27%
S	20.31%	20.16%

C. Methyl-2-chloro-6-fluoro-N-[(3,4,5-trichloro-2-thiophene)carbonyl]-benzenecarboximidothioate.



2-Chloro-6-fluorobenzthioimide methylsulfate salt (35.5 g, 0.11 mol) and 155 mL of toluene were placed into a 1 L three-necked roundbottom flask with a bottom drain and overhead stirrer. The solution was cooled with an ice bath, and to this slurry was added triethylamine (32 mL, 23.2 g, 0.23 mol). A mild exotherm was noted. A solution of 3,4,5-trichloro-2-thiophenecarbonyl chloride (30.69 g, 0.12 mol) was added dropwise over 30 min at 0°C. After 1 h of cooling, the ice bath was removed and the mixture was stirred at 25°C for 3 days. The solution was washed with two 250 mL portions of water, and the toluene was evaporated to give 41.3 g of tan solid. This solid was washed with CH₃CN, filtered and dried to give 32.8 g of white solid, m.p. 124-126°C, for a weight % yield of 70.2 %.


A purified sample of the adduct was obtained by recrystallization from warm CH₃CN followed by vacuum oven drying at 60°C for 3 h; m.p. 125-128°C. Elemental analysis results are summarized in the table below.

	<u>Theoretical</u>	<u>Found</u>
C	37.43 %	37.18 %
H	1.45 %	1.43 %
N	3.36 %	3.62 %
Cl	34.00 %	34.08 %
F	4.55 %	5.30 %
S	15.37 %	15.64 %

D. 1-Methyl-3-(2-chloro-6-fluorophenyl)-5-(3,4,5-trichloro-2-thienyl)-1H-1,2,4-triazole.

The adduct intermediate of part C (6.1 g, 0.0146 mol) and 75 mL toluene were placed into a 250 mL three-necked flask and brought to reflux. Methylhydrazine (1.6 mL, 1.39 g, 0.0301 mol) as a solution in 25 mL of toluene was added dropwise to the refluxing solution over a period of 1 h. The solution was refluxed for an additional 2 h, then was cooled to 25°C, washed with aqueous Na₂CO₃, then water, and the organic phase was evaporated to give a viscous oily residue. The ratio of regio isomers by GC area % analysis was 13.7/1 in favor of the desired isomer. The oily residue was triturated with a small volume of CH₃CN, cooled to 0°C in a freezer for 1 h, and the resulting white crystals were collected on a buchner funnel and air-suction dried to give 4.1 g (70%) of the desired product, m.p. 128-130.5°C, with a GC area % purity of 98.3 % and containing 1.4 % of a regio isomer impurity.

A preferred synthetic method is illustrated in the following Scheme I:




The first step of this process involves preparation of the thioamide starting material. The thioamide was converted to the thioimide using dimethyl sulfate in 1,4-dioxane. Any known imide forming procedure known in the literature can be used for this transformation. Common methylating agents such as methyl iodide, methyl bromide and dimethyl sulfate can be used. Any common solvent compatible with the reaction conditions can be used, with toluene, acetonitrile, 1,4-dioxane, THF, and 1,2-dichloroethane most convenient. Reaction temperatures range from RT to the reflux temperature of the solvent. The thioimide can be isolated as its salt or used directly without isolation in the next transformation.

The thioimide was next acylated with 3-methyl-2-thiophenecarbonyl chloride ("acid chloride") to give the acyl thioimide adduct. Any known acylation conditions can be used for this transformation. Any common organic and inorganic base can be used, with Na_2CO_3 , NaHCO_3 , pyridine and triethylamine most convenient. Solvents preferred include 1,4-dioxane, THF, dichloromethane, and 1,2-dichloroethane, but any solvent compatible with the

reaction conditions can be used. Reaction temperatures in the 0° to 60°C range are suitable, with temperatures near RT most convenient. The acyl thioimide was isolated by dilution of the reaction mixture with water and filtration followed by air drying. This acyl thioimide was of sufficient purity to be used directly in the following cyclization step.

The acyl thioimide was cyclized to the 1,2,4-triazole ring system by treatment with methylhydrazine. The methylhydrazine could be added neat or as a solution in a compatible solvent such as water. Any solvent compatible with the reaction conditions can be used, with toluene, 1,4-dioxane, THF, and short chain alcohols preferred. The methylhydrazine can be added all at once to the reaction mixture, or added in portions over a 1 hour time period. The cyclization can be carried out in the temperature range of RT to reflux temperature of the solvent being used. Ratios of the intermediate **3** to its off-isomer **4** range from 6:1 to 40:1 depending upon the reaction conditions used.



It was found most convenient to use 1,4-dioxane at a cyclization temperature of 80°C to give a 30:1 ratio of isomers. Workup consisted of removal of the solvent followed by crystallization from a suitable solvent such as 95% EtOH. Alternately, the reaction mixture could be diluted with water and filtered to provide the intermediate **3**. Air drying provided product of sufficient purity for the next bromination step.


The last two steps illustrated in Scheme I are the bromination of triazole **3** to give the dibromo

intermediate **2**, followed by removal of one of the bromine atoms. Any standard brominating reagent known in the literature can be used, with Br₂ being the most convenient. Two to five molar equivalents of Br₂ can be
5 used at temperatures from 25° to reflux temperature of the solvent. Time of reaction range from 1 hour to 24 hours. Any solvent compatible with bromination conditions can be used such as 1,4-dioxane, 1,2-dichloroethane, and acetic acid. The generated HBr can
10 be neutralized by running the reaction in the presence of a proton acceptor such as sodium acetate. It was found most convenient to run the reaction in acetic acid with sodium acetate using four equivalents of Br₂.

The dibromo analogue **2** can be isolated or one may
15 proceed directly into the next chemical transformation, if desired, without isolation. In the last step, zinc dust is added to the reaction mixture to reduce off the 5-bromine on the thiophene ring. Any known methods of aromatic halogen reduction could be used, but zinc dust
20 was found most convenient. Two to three equivalents of zinc can be used; the extra molar equivalents were needed to reduce unreacted Br₂. The temperature of the reduction ranged from 25° to 90°C. This reduction is highly selective and leaves the 4-bromine of the thiophene ring
25 unaffected. The product is conveniently isolated by dilution of the reaction mixture with water followed by filtration.

Another preferred synthetic method is illustrated in the following Scheme II:

Scheme II




- Conditions for each step in Scheme II are as follows: a) $\text{H}_2\text{S}/\text{Et}_3\text{N}/\text{Pyridine}$, -20°C , b) $(\text{CH}_3\text{O})_2\text{SO}_2$, 1,4-dioxane, 80°C ,
 5 c) pyridine, 4-bromo-TAC, 30°C ,
 d) $\text{MeNHNH}_2/\text{H}_2\text{O}$, 1,4-Dioxane, 80°C .

The 4-bromo-3-methyl-2-thiophenecarbonyl chloride used in step **c** of Scheme I can be prepared using the procedure illustrated in Scheme III:

10

Scheme III



- Conditions for each step in Scheme III are as follows:
 e) $\text{Br}_2/\text{succinimide}$, f) 10% Pd/C , DPPP, 0.2 mol %, CO

pressure, EtOH, NaOAc, g) i) 2 Br₂/2 NaOAc/HOAc, 80°C,
ii) Zn dust, HOAc/H₂O, h) NaOH/H₂O, i) SOCl₂/DMF, 1,2-DCE.


Scheme III illustrates use of ethyl 3-methyl-2-thiophenecarboxylate as the intermediate utilized in step
5 **g**, but any short chain alkylester of 3-methyl-2-thiophenecarboxylic acid may be used. These include, but are not limited to, methyl, ethyl, propyl or butyl. Most convenient are methyl or ethyl esters of 3-methyl-2-thiophenecarboxylic acid.

10 One aspect of the invention is a novel preparation of ethyl-3-methyl-2-thiophenecarboxylate via a palladium-catalyzed carboxylation of 2-bromo-3-methylthiophene.

Methyl 3-methyl-2-thiophenecarboxylate may be prepared by a Grignard reaction of 2-bromo-3-
15 methylthiophene with dimethylcarbonate. Either the methyl or ethyl ester of 3-methyl-2-thiophene carboxylic acid may be prepared by Fisher esterification with the appropriate alcohol, or by reaction of 3-methyl-2-thiophenecarboxylic acid chloride with the appropriate
20 alcohol.

Yet another preferred embodiment is illustrated in Scheme IV:

Scheme IV



Noteworthy features of the embodiment illustrated in
 5 Scheme IV include:

- 1) the methyl sulfate salt of the benzthioimide is used;
- 2) 1,2-dichloroethane is used as the solvent in each
 step;
- 3) the synthesis is carried out in one pot, without
 10 isolation of intermediates;
- 4) octane is added in the final step to promote
 crystallization of the product.

The individual steps in the Scheme IV are described
 in detail hereafter:

- 15 The first step of this process involves the
 conversion of 2-chloro-6-fluorobenzonitrile to the
 thioamide. Any of the methods known in the chemical

literature can be used for this thioamide formation reaction. Sodium sulfide can be used as the sulfur source, but it has been found most convenient to use hydrogen sulfide gas. Reaction temperatures used are in
5 the -35° to 50° C range, with -10° to RT most convenient. Any common solvent compatible with the reaction conditions can be used. Pyridine and ethanol are suitable. Any common amine base, for example triethylamine, can be used.

10 The next three distinct chemical transformations are most conveniently carried out in one process step, without isolation of two intermediates. The two intermediates can be isolated and characterized if desired. In the first transformation, the 2-chloro-6-
15 fluorothiobenzamide is converted to the methyl thioimide. Any known imide forming procedure known in the literature can be used for this transformation. Common methylating agents such as methyl iodide, methylbromide and dimethyl sulfate can be used. Any
20 common solvent compatible with the reaction conditions can be used, with toluene, acetonitrile, THF, and 1,2-dichloroethane most convenient. Reaction temperatures range from RT to the reflux temperature of the solvent. The thioimide can be isolated as its salt or used
25 directly without isolation in the next transformation.

The thioimide is next acylated with 3,4,5-trichloro-2-thiophenecarbonyl chloride to give the acylated adduct. This adduct can be isolated and characterized if desired, but was found to be most
30 conveniently used directly without isolation. Any known acylation conditions can be used for this transformation. Any common organic and inorganic base can be used, with Na_2CO_3 , NaHCO_3 , pyridine and triethylamine most convenient. Solvents preferred include THF,
35 dichloromethane, and 1,2-dichloroethane, but any solvent

compatible with the reaction conditions can be used. Reaction temperatures in the 0° to 60° C range are suitable, with temperatures near RT most convenient.

The acylated adduct is finally cyclized to the 1,2,4-triazole ring system by treatment with methylhydrazine. A two step procedure using hydrazine to give the unsubstituted triazole followed by methylation could be used, but it is convenient to use methylhydrazine directly. The methylhydrazine can be added neat or as a solution in a compatible solvent such as water. Any solvent compatible with the reaction conditions can be used, with toluene, THF, and 1,2-dichloroethane preferred. The methylhydrazine can be added all at once to the reaction mixture, or added in portions over a 1 hour time period. The cyclization can be carried out in the temperature range of RT to reflux temperature of the solvent being used. Ratios of the desired product to its off-isomer range from 1:4 to 50:1 depending upon the reaction conditions used. It is most convenient to use 1,2-dichloroethane at a cyclization temperature of 70° C to give a 38:1 ratio of 1-methyl-3-(2-chloro-6-fluoro)phenyl-5-(3,4,5-trichloro-2-thienyl)-1H-1,2,4-triazole to its "off isomer."

Example 4

This example illustrates the process of Scheme IV.

A. Preparation of 2-chloro-6-fluorobenzthioamide in pyridine/triethylamine solvent.

2-Chloro-6-fluorobenzonitrile, (99.1%, 62.2 g, 0.40 mol) was weighed into a 1-L three-necked roundbottom flask equipped with a condenser and an overhead electric stirrer, along with Et₃N (78 mL, 56.6 g, 0.56 mol) and 180 mL (176.04 g, 2.23 mol) of pyridine. The reactor was purged with a slow stream of N₂ and vented to a 13% bleach solution. The stirring solution was cooled to -19°C in a

CCl₄/dry ice bath, and H₂S gas (33.6 g, 0.99 mol) was sparged below the liquid surface at a rate of 0.4 g/min over a period of 82 min. During the gas addition, the solution temperature rose to -11°C. The yellow-green
5 solution was allowed to gradually warm to 25°C and stir overnight with a slow N₂ purge of the reactor head space into bleach solution. The solution was poured into 1.6 L of ice water, stirred, and the resulting white crystals were collected on a buchner funnel and rinsed with
10 additional water. After 2 h of air drying, the moist filter cake was vacuum oven dried for 5 h at 65°C to give 54.5 g of 2-Chloro-6-Fluorobenzthioamide (72% wt. % yield), mp 155-160°C, having a GC area % purity of 98.8 % and containing 1.2 % 2-Chloro-6-fluorobenzonitrile.

15 B. Preparation of 2-chloro-6-fluorobenzthioamide in ethanol/triethylamine solvent.

2-Chloro-6-fluorobenzonitrile, (99.1%, 15.2 g, 0.10 mol) was weighed into a 1-L three-necked roundbottom flask equipped with a condenser and an overhead electric
20 stirrer, along with Et₃N (41 mL, 29.8 g, 0.29 mol) and 54 mL of 95% ethanol. The solution was cooled to 0°C with an ice bath and H₂S gas (13.5 g, 0.40 mol) was sparged below the liquid surface at a rate of 0.2 g/min over a period of 70 min. The ice bath cooling was continued for
25 2 h, then the flask was allowed to gradually warm to room temperature. The yellow solution was poured into 426 g of stirring ice water in a 1 L Erlenmeyer flask, and the resulting white solid was collected on a buchner funnel and rinsed with additional water. This filter cake was
30 dissolved in 1,2-dichloroethane and evaporated to dryness on a rotary evaporator (67°C bath) to ensure dryness, giving 13.42 g (72 % weight % yield) of the thioamide as a white solid.

C. 3,4,5-Trichloro-2-thiophenecarbonyl chloride

A mixture of 20.4 g (0.088 mole) of 3,4,5-trichloro-2-thiophenecarboxylic acid, 7.3 mL (0.1 mole) of thionyl chloride, 0.2 mL of DMF and 80 mL of 1,2-dichloroethane was heated at reflux temperature for 3 h. The reaction mixture became a clear solution after 1 h of heating. The mixture was allowed to cool to RT, concentrated *in vacuo* and exposed to high vacuum to give 22.0 g (>98 % wt) an oil which solidified upon standing to an off-white powder, mp 37°-41° C.

10 D. 1-Methyl-3-(2-chloro-6-fluorophenyl)-5-(3,4,5-trichloro-2-thienyl)-1H-1,2,4-triazole

A mixture of 94.82 g (0.5 mole) of 2-chloro-6-fluorobenzthioamide, 50 mL (0.525 mole) of dimethyl sulfate, and 800 mL of 1,2-dichloroethane was heated at reflux temperature for 1 h. LC analysis indicated complete conversion to the thioimide. The reaction solution was cooled to RT with ice-bath cooling, and the thioimide salt precipitated from the solution. To the thick slurry was added 100 mL (1.25 mole) of pyridine. The salts instantly dissolved into solution, followed by immediate precipitation of pyridine salts. After further cooling to 10° C, 125 g (0.5 mole) of 3,4,5-trichloro-2-thiophenecarbonyl chloride dissolved in 200 mL of 1,2-dichloroethane was added in three portions. Temperature of the reaction mixture rose to 20° C during each addition and was allowed to cool back to 10° C before the next addition. The reaction mixture was allowed to stir at RT for 0.5 h. LC analysis indicated major acylated thioimide and <2% area of starting thioimide. To the slurry was added 1.0 L of water and the layers separated. LC internal standard analysis of the organics indicated a 94% in-pot yield of the acylated intermediate. The 1,2-dichloroethane solution was heated to 70° C, and a solution of 40 mL (0.75 mole) of methylhydrazine in 60 mL of water was added via a pump over 20 min. The

temperature of the reaction mixture was maintained between 70° and 73° C. The solution was heated an additional 2 h, when GC analysis indicate no remaining acylated intermediate and a 38:1 ratio of desired product/off-isomer. The reaction mixture was cooled to RT, and 500 mL of octane followed by 1 L of 0.5 N LiOH was added. After stirring for 10 min, the layers were separated. The organic solution was then stirred with 750 mL of a 2% bleach solution for 0.25 h to remove remaining mercaptan compounds. The temperature of the mixture rose to 31° C during this reaction. The layers were separated and upon concentration of the 1,2-dichloroethane on a roto-evaporator, a tan sandy precipitate formed. This solid was filtered and exposed to vacuum oven drying at 40° C to give 144.8 g (73% wt) of the desired product as a tan crystalline powder. GC analysis indicated <0.2% area of the off-isomer, and LC internal standard analysis indicated a 98% purity, giving an overall isolated 72% yield of 1-methyl-3-(2-chloro-6-fluoro)phenyl-5-(3,4,5-trichloro-2-thienyl)-1H-1,2,4-triazole.

Example 5

3-(2-Chloro-6-fluorophenyl)-1-methyl-5-(3-methyl-4-bromo-2-thienyl)-1H-1,2,4-triazole

This example illustrates the process of Scheme I.

A. 2-Chloro-6-fluorobenzenecarbothioamide

2-Chloro-6-fluorobenzonitrile (1230 g, 7.91 mol) was weighed into a 22 L jacketed glass reactor (without bottom drain), along with Et₃N (1.5 L, 10.76 mol) and pyridine (2.4 L, 29.67 mol) and the stirring solution was cooled to -18°C under a blanket of N₂. The reactor N₂ purge mineral oil bubbler was vented to a carboy containing 16 L of 12% bleach solution. Hydrogen sulfide gas (407 g, 11.94 mol) was introduced below the surface of the cold reactor solution over a period of 5.5 hrs.

During this time the reactor temperature rose from -18 to -4°C. The solution was stirred overnight at -7°C (16 hrs), sampled by GC to ensure complete conversion of 2,6-CFBN, then quenched by vacuum transfer of the reactor solution into a second 22 L flask containing 14 L of cold water stirring at 300 RPM (Note: Quenching into a well-stirred tank of cold water produces small white particles of easily filtered product). After the vacuum transfer was completed, another 2 L of cold water was added to the stirring slurry. The light yellow slurry of white solid was drained from the quench tank into bottles, and the solid product was collected by filtration through a course glass buchner funnel, followed by washing of the solid product with an excess of de ionized water. The white solid was air suction dried for 3 h, air dried in a glass pan overnight, then dried in a vacuum oven at 50°C at 0.3 mm Hg for 8 hrs to give 1004 g (67%) of dry 2-chloro-6-fluorobenzenecarbothioamide, m.p. 153-157°C.

B. Methyl 2-chloro-6 fluoro-N-[(3-methyl-2-thiophene)carbonyl]benzenecarboximidothioate
To a 22 L jacketed glass reactor under a N₂ purge was added 6 L of 1,4-dioxane via a pump. After the mechanical stirrer had been started, 1140 g (6.0 mol) of 2-chloro-6 fluorobenzenecarbothioamide was added to the reactor, followed by 630 mL (6.65 mol) of dimethyl sulfate. The reaction mixture was heated to 80°C for 1.5 h. During the course of heating, the reaction mixture became a solution near 55°C, with a precipitate reforming as the reaction proceeded. LC analysis indicated <2% area starting thioamide and 96% area methyl 2-chloro-6-fluorobenzenecarboximidothioate methylsulfonate salt ("thioimide"). The reaction mixture was cooled to 30°C, and 1.2 L (15 mol) of pyridine was added. When the reaction temperature had fallen back below 30°C, 970 g (6.04 mole) of 3-methyl-2-thiophenecarbonyl chloride

("acid chloride") was added via a PE dropping funnel over 20 min. An exotherm of seven degrees was observed during this addition. After stirring at room temperature for 1 h, LC analysis indicated < 3% area of the thioimide and 93% area acyl thioimide. To the reaction mixture was added 9 L of water, and the mixture stirred at room temperature for 1 h. The mixture was filtered through a crock filter with vacuum removal of the mother liquor into a second 22 L glass reactor. The white solid was air dried in a glass pan overnight to give 2000 g of a white powder, 98% LC area. This material was used directly in the next step without further purification.

C. 3-(2-Chloro-6-fluorophenyl)-1-methyl-5-(3-methyl-2-thienyl)-1H-1,2,4-triazole

To a 22 L jacketed glass reactor under a N₂ purge was added 9 L of 1,4-dioxane via a pump. The N₂ purge outlet was connected to a carboy containing 12 L of fresh 12% wt bleach solution. To the stirred dioxane was added 1970 g (6 mol) of methyl 2-chloro-6 fluoro-N-[(3-methyl-2-thiophene)carbonyl]benzenecarboximido-thioate and the mixture heated to 80°C. During heating the mixture became a clear solution. Upon reaching 80°C, a solution of 400 mL (7.5 mol) of methylhydrazine in 600 mL of water was added over 20 min via a pump. The mixture was heated near 80°C for 3.5 h, when GC area analysis indicated 2% starting acyl thioimide, 94% desired product, and 3% off-isomer 4. The reaction mixture was cooled to 30°C and vacuum transferred to a roto-evaporator connected to bleach trapping. The reaction mixture was concentrated to an amber oil, dissolved in 800 mL of 95% EtOH, and transferred to a 4 L Erlynmeyer flask. The solution was scratched and seeded with the title compound and allowed to crystallize in a refrigerator overnight. After filtration the precipitate was air dried in a glass pan

overnight to give 1376 g of an off-white powder, 99% area GC. This material was used directly in the next step without further purification.

D. 3-(2-Chloro-6-fluorophenyl)-1-methyl-5-(3-methyl-4-bromo-2-thienyl)-1H-1,2,4-triazole

The triazole 3-(2-Chloro-6-fluorophenyl)-1-methyl-5-(3-methyl-2-thienyl)-1H-1,2,4-triazole (1715 g, 5.57 mol) and anhydrous sodium acetate (1830 g, 22.3 mol) were weighed as solids into a 22 L jacketed glass reactor under a slow N₂ purge, followed by 8.5 L glacial AcOH. Heating of the mixture was begun, and at 66°C a solution of Br₂ (1.15 L, 22.3 mol) in 2.5 L glacial AcOH was added by peristaltic pump over a period of 40 min., at a rate to keep the reaction temperature below 85°C. During heat up most of the solids dissolve. The stirring red solution was heated 1 h at 80°C and then a sample was removed and analyzed by GC to verify >99% conversion to the dibromo intermediate. Cooling was placed on the jacket, and 1695 g ice was added to the solution, which dropped the temperature from 70°C to 50°C. At 38°C zinc powder (815 g, 12.5 mol) was added in 55 g shots over a period of 40 min to dispel excess Br₂. During this addition the reactor temperature rose to 71°C. Additional zinc powder (650 g, 9.9 mol) was then added in 50 g shots over a period of 50 min with a stream of N₂ passing through the reactor to dispel H₂ gas safely. During the addition of this second portion of zinc powder, jacket heating (85°C) was applied to bring the reaction mixture to 78°C (Note: zinc addition at this point leads to H₂ gas evolution and should be done at such a rate as to control the rate of evolution and maintain precautions against flammability and explosion hazards). After zinc addition was completed, the reactor was stirred at 78°C for 15 min, then was sampled for GC analysis (Note: the reactor can be maintained at this

point at 80°C overnight to keep the product dissolved if insufficient time exists to carry out the filtration). The solution was filtered warm through a layer of celite on a course glass buchner funnel (pre warmed at 100°C) to
5 remove residual zinc particulates. The celite cake was washed with a small amount of additional glacial AcOH. The clear light yellow filtrate was diluted with cold water while stirring until the cloud point was reached and product crystallization began. Excess cold water was
10 then added to precipitate the remainder of the solid product. Part of this dilution was done in 4 L Erlenmeyer flasks with magnetic stirring and part was done in a second 22 L reactor. Approximately eight 2 L portions of solution were filtered from the reactor, and
15 approximately 18 L of quench water was used. The white solid product was collected on course glass buchner funnels, rinsed with additional de-ionized water, air suction dried for a few hours, then transferred to glass trays and air dried overnight. A final vacuum oven
20 drying for 2.5 hrs at 0.3 mm Hg and 30°C was carried out to give 2236 g of 3-(2-Chloro-6-fluorophenyl)-1-methyl-5-(3-methyl-4-bromo-2-thienyl)-1H-1,2,4-triazole as a white solid (theory = 2154 g), mp 118-120°C. A GC area % analysis indicated a purity of 98.1%.

25

Example 64-Bromo-3-methyl-2-thiophenecarbonyl chloride

This example illustrates the process of Scheme III.

A. Ethyl 3-methyl-2-thiophenecarboxylate via palladium catalyzed carbonylation of 2-bromo-3-methylthiophene

30 (Scheme III, step f)

2-Bromo-3-methylthiophene (Lancaster, 26.4 g, 0.149 mol), Pd(OAc)₂ (0.213 g, 0.95 mmol), 1,3-bis(diphenylphosphino) propane (0.49 g, 1.2 mmol), Na₂CO₃ (21.1 g, 0.20 mol), and 200 mL abs EtOH were placed into

a 450 mL Hastalloy-C stirred pressure reactor, purged with N₂, then pressurized with 490 psig CO, and the stirring reactor was heated to 120°C for 36 hrs. Analysis of the reaction mixture by gas chromatography showed 1.3 % and 96.9 % area % starting bromomethylthiophene and carbonylation product, respectively. After cooling and venting, the reactor contents were filtered and evaporated to give a light yellow oil/solid mixture. This residue was extracted with 1,2-dichloroethane and water, and the organic phase was evaporated to give an oil. Filtration of this oil to remove residual salt gave 20.7 g (81% wt. % yield) of orange oil as ethyl 3-methyl-2-thiophenecarboxylate. ¹³C NMR {¹H} CDCl₃: δ 162.8, 145.9, 131.7, 129.9, 127.0, 60.6, 15.9, 14.4 ppm.

B. Ethyl 4-bromo-3-methyl-2-thiophenecarboxylate (Scheme III, step g)

A solution of ethyl 3-methyl-2-thiophenecarboxylate (20.0 g, 0.118 mol) and sodium hydroxide (12.3 g, 0.307 mol) in acetic acid (75 mL) was heated to 60 °C. Bromine (46.9 g, 0.294 mol) was added dropwise at such a rate so as to maintain the temperature of the reaction mixture at <85 °C. When the addition was complete, the resulting solution was stirred at 85 °C for 6 hours, at which time analysis by gas chromatography/mass spectrometry showed complete conversion to ethyl 4,5-dibromo-3-methyl-2-thiophenecarboxylate. The solution was allowed to cool to 50 °C and zinc dust (15.4 g, 0.236 mol) was added in 3 gram portions such that the exotherm was controlled to remain below 85 °C. When the addition was completed, the resulting slurry was stirred at 85 °C for one hour. The solution was filtered hot through a small bed of celite. Water (300 mL) was added and the mixture was extracted with heptanes (300 mL). The organic phase was washed with water, then concentrated to dryness to give 26.9 g (89%) as an off white oil which slowly crystallized upon

standing at ambient temperature. In a similar fashion, methyl 3-methyl-2-thiophenecarboxylate may be converted methyl-4bromo-3-methyl-2-thiophenecarboxylate in 97% yield.

5 C. 4-Bromo-3-methyl-2-thiophenecarboxylic acid
(Scheme III, step h)

To a solution of ethyl 4-bromo-3-methyl-2-thiophenecarboxylate (5.0 g, 0.0201 mol) in THF/MeOH/H₂O (4:1:1, v/v/v, 50 mL) was added NaOH (1.00 g, 0.0251 mol)
10 and the resulting mixture was stirred at ambient temperature overnight. The mixture was acidified by adding 6 N HCl (100 mL) and water (100 mL). The resulting fine white precipitate was filtered off, washed with water and dried to give 3.80 g (86%) of 4-bromo-3-methyl-2-thiophenecarboxylic acid as a fine white solid,
15 mp 188-189°C.

D. 4-Bromo-3-methyl-2-thiophenecarbonyl chloride (Scheme III, step i)

A slurry of 1.11 g (5 mmole) of 4-bromo-3-methyl-2-thiophenecarboxylic acid, 0.44 mL (6 mmole) of thionyl chloride, 5 drops of DMF, and 10 mL of 1,2-dichloroethane was heated at 80° C for 1.5 h. The clear solution was cooled and concentrated on a roto-evaporator. The residue was dissolved in 5 mL of 1,2-dichloroethane and
25 reconcentrated on a roto-evaporator to give 1.2 g of a tan solid, 97 % pure (GC area). This acid chloride was used without further purification.

Example 7

30 3-(2-Chloro-6-fluorophenyl)-1-methyl-5-(4-bromo-3-methyl-2-thienyl)-1H-1,2,4-triazole

This example illustrates the process of Scheme II.

A. Methyl 2-chloro-6-fluoro-N-[(4-bromo-3-methyl-2-thiophene)carbonyl]benzenecarboximidothioate (Scheme II, step c)


A mixture of 0.95 g (5 mmole) of 2-chloro-6-fluorobenzenecarbothioamide, 0.5 mL (5.5 mmole) of dimethyl sulfate, and 10 mL of 1,4-dioxane was heated at 80° C for 1 h. The clear solution was allowed to cool to room temperature, where a white precipitate formed. To the slurry was added 1.0 mL (12.5 mmole) of pyridine, followed by 1.2 g (5 mmole) of 4-bromo-3-methyl-2-thiophenecarbonyl chloride. The reaction mixture turned brown in color and a gummy insoluble material formed which made stirring difficult. The mixture was heated at 50° C for 1.5 h. The mixture was poured onto ice and a gummy solid formed which solidified upon stirring. Stirring was continued for 1 h, and the mixture was filtered to give 1.74 g (85% wt) of a tan powder, mp 106°-109°. This material was used without further purification.

B. 3-(2-Chloro-6-fluorophenyl)-1-methyl-5-(4-bromo-3-methyl-2-thienyl)-1H-1,2,4-triazole (Scheme II, step d)

A mixture of 1.22 g (3 mmole) of methyl 2-chloro-6-fluoro-N-[(4-bromo-3-methyl-2-thiophene)carbonyl]-benzenecarboximidothioate in 5 mL of 1,4-dioxane was heated to 75° C. To the solution was added in one portion 0.24 mL (4.5 mmole) of methylhydrazine in 0.5 mL of water. After heating here for 1.5 h, the solution was cooled and poured onto ice. The initially formed gummy solid was stirred for 1 h and filtered to give 1.03 g (89% wt) of an off-white powder. GC analysis indicated 96.4/1.7 area ratio of the desired product to off-isomer.

Claims

1. A process for preparing a compound of the formula (1)




5 wherein

Ar is substituted phenyl;

Y is O or S;

R³ is selected from H, halo, lower alkyl, (C₇-C₂₁) straight or branched chain alkyl, hydroxy, lower alkoxy, haloalkyl, haloalkoxy, alkoxyalkyl, alkoxyalkoxy, lower alkenyl, lower alkynyl, haloalkenyl, CN, NO₂, COR⁶, CO₂R⁶, CON(R⁶)₂, (C₃-C₆) cycloalkyl, S(O)_mR⁶, -OSO₂R⁶, SCN, - (CH₂)_nR⁶, -CH=CHR⁶, -C≡CR⁶, -(CH₂)_qOR⁶, -(CH₂)_qSR⁶, - (CH₂)_qNR⁶R⁶, -O(CH₂)_qR⁶, -S(CH₂)_qR⁶, -NR⁶(CH₂)_qR⁶,



, pyridyl, substituted pyridyl, isoxazolyl, substituted isoxazolyl, naphthyl, substituted naphthyl, phenyl, substituted phenyl, thienyl, substituted thienyl, pyrimidyl, substituted pyrimidyl, pyrazolyl, or substituted pyrazolyl;

R⁴ and R⁵ are independently H, halo, lower alkyl, lower alkoxy, haloalkyl, haloalkoxy, CN, CO₂R⁶, CON(R⁶)₂, or S(O)_m alkyl, or

if R^4 and R^5 are attached to adjacent carbon atoms, they may join to form a 5 or 6 member saturated or unsaturated carbocyclic ring which may be substituted by 1 or 2 halo, lower alkyl, lower alkoxy or haloalkyl groups;

R^6 is H, lower alkyl, haloalkyl, lower alkenyl, lower alkynyl, phenyl, or substituted phenyl;

m is 0, 1, or 2;


n is 1 or 2;

10 p is an integer from 2 to 6; and


q is 0 or 1;

which comprises the steps of:


(a) reacting a compound of formula (2)



15 wherein Ar is as defined above and R^7 is lower alkyl, or an acid addition salt thereof, with an acid chloride of the formula (3)



wherein Y, R^3 , R^4 , and R^5 are as defined in formula (1),
 20 in an inert organic solvent the presence of an organic or inorganic base to produce an adduct-intermediate of formula (4)




wherein Ar, Y, R³, R⁴, and R⁵ are as defined above and R⁷ is lower alkyl; and

(b) with or without prior isolation of the adduct-intermediate of formula (4), reacting said adduct-intermediate with methyl hydrazine to produce the compound of formula (1).


2. A process of claim 1 wherein an acid addition salt of the reactant of formula (2) is used.

3. A process of claim 2 wherein the acid addition salt of the reactant of formula (2) is the hydroiodide or the methyl sulfate salt.


4. A process of claim 1 wherein a compound of formula (2a)




wherein R¹ and R² are independently F or Cl, and R⁷ is lower alkyl, or an acid addition salt thereof, is reacted with an acid chloride of the formula (3a)



wherein R³, R⁴ and R⁵ are independently H, CH₃, Cl, or Br to produce the adduct-intermediate of formula (4b)



wherein R^1 , R^2 , R^3 , R^4 , R^5 , and R^7 are as defined above,
 and, with or without isolation of said adduct-
 intermediate of formula (4b), said adduct-intermediate is
 reacted with methyl hydrazine to produce the compound of
 5 formula (1b)



wherein R^1 , R^2 , R^3 , R^4 , and R^5 are as defined above.


5. A process of claim 4 wherein R^3 and R^7 are CH_3 ,
 and R^4 and R^5 are both H or both Br, or R^4 is Br and R^5 is
 10 H.

6. A process of claim 5 wherein R^1 is F, R^2 is Cl,
 R^4 is Br, and R^5 is H or Br.


7. A process of claim 6 wherein

R^5 is Br, and

15 the product of formula




is reacted with zinc dust to provide a compound of
 formula



8. A process of any one of claims 1 to 7 wherein the adduct intermediate is reacted with methyl hydrazine in 1,4-dioxane.

9. A compound of the formula



5


wherein

Ar is substituted phenyl,

Y is O or S,

R³ is selected from H, halo, lower alkyl, (C₇-C₂₁)

10 straight or branched chain alkyl, hydroxy, lower alkoxy, haloalkyl, haloalkoxy, alkoxyalkyl, alkoxyalkoxy, lower alkenyl, lower alkynyl, haloalkenyl, CN, NO₂, COR⁶, CO₂R⁶, CON(R⁶)₂, (C₃-C₆) cycloalkyl, S(O)_mR⁶, -OSO₂R⁶, SCN, - (CH₂)_nR⁶, -CH=CHR⁶, -C≡CR⁶, -(CH₂)_qOR⁶, -(CH₂)_qSR⁶, -
15 (CH₂)_qNR⁶R⁶, -O(CH₂)_qR⁶, -S(CH₂)_qR⁶, -NR⁶(CH₂)_qR⁶,



-NR⁶-C(=O)-N(R⁶), -NR⁶-C(=O)-OR⁶, or -S-C(=O)-R⁶, pyridyl, substituted pyridyl, isoxazolyl, substituted isoxazolyl, naphthyl, substituted naphthyl, phenyl, substituted
20 phenyl, thienyl, substituted thienyl, pyrimidyl, substituted pyrimidyl, pyrazolyl, or substituted pyrazolyl;

R⁴ and R⁵ are independently H, halo, lower alkyl, lower alkoxy, haloalkyl, haloalkoxy, CN, CO₂R⁶, CON(R⁶)₂, or S(O)_m alkyl, or

if R⁴ and R⁵ are attached to adjacent carbon atoms, they may join to form a 5 or 6 member saturated or unsaturated carbocyclic ring which may be substituted by 1 or 2 halo, lower alkyl, lower alkoxy or haloalkyl groups;

R⁶ is H, lower alkyl, haloalkyl, lower alkenyl, lower alkynyl, phenyl, or substituted phenyl;

m is 0, 1, or 2;


n is 1 or 2;

p is an integer from 2 to 6;

q is 0 or 1; and

R⁷ is lower alkyl.

10. A compound of claim 9 having the formula (4a)



wherein

R¹ and R² are independently F or Cl;

R³, R⁴, and R⁵ are independently H, CH₃, Cl or Br;

and

R⁷ is lower alkyl.

11. A compound of claim 10 wherein


R⁷ is methyl, R¹ is F, R² is Cl, and

(a) R³, R⁴, and R⁵ are each Cl;

(b) R³ and R⁴ are each Br and R⁵ is H; or

(c) R^3 is CH_3 , R^4 is Cl or Br, and R^5 is H.

12. The methyl sulfate salt of the compound of the formula



INTERNATIONAL SEARCH REPORT

Int. l. Application No

PCT/US 99/24860

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D409/04 C07D405/04 C07D333/38 C07D307/68 C07C327/58

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 609 459 A (KUMIAI CHEMICAL INDUSTRY CO. LTD.) 10 August 1994 (1994-08-10) page 12, line 1 -page 13, line 13; claims	1-12
A	BE 786 562 A (GRUPPO LEPITATIT S.P.A.) 16 November 1972 (1972-11-16) page 2, line 29 -page 3, line 20; claims	1-12
A	T.W. STROHMEYER ET. AL.: "New Synthesis of 2,4-Dialkyl(or diaryl) pyrazolo-1,5-a! 1,3,5-triazines." JOURNAL OF HETEROCYCLIC CHEMISTRY, vol. 22, no. 1, January 1985 (1985-01), pages 7-10, XP000881466 page 8; table 1	9-12

	---/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

6 March 2000

Date of mailing of the international search report

23/03/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Helps, I

INTERNATIONAL SEARCH REPORT

Int. l. Application No

PCT/US 99/24860

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	M. A. PEREZ ET. AL. : "Regioselective Synthesis of 1,2,4-Triazole and 1,2,4-Oxadiazole Derivatives." SYNTHESIS, vol. 1983, no. 6, June 1983 (1983-06), pages 483-6, XP002132293 cited in the application whole article	1-12
P,A	WO 98 47894 A (DOW AGROSCIENCES LLC) 29 October 1998 (1998-10-29) claims; examples	1-12

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/24860

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 609459	A	10-08-1994	JP 6041093 A	15-02-1994
			JP 6041094 A	15-02-1994
			JP 6056801 A	01-03-1994
			US 5466705 A	14-11-1995
			ES 2095657 T	16-02-1997
			WO 9402471 A	03-02-1994
BE 786562	A	16-11-1972	AR 195492 A	15-10-1973
			AR 194297 A	29-06-1973
			AR 194298 A	29-06-1973
			AR 200485 A	15-11-1974
			AT 317210 B	26-08-1974
			AT 319236 B	10-12-1974
			AU 461487 B	29-05-1975
			AU 4324672 A	13-12-1973
			CA 1015361 A	09-08-1977
			CH 558372 A	31-01-1975
			CH 559195 A	28-02-1975
			CH 560707 A	15-04-1975
			DD 101574 A	12-11-1973
			DE 2235544 A	15-03-1973
			DE 2265212 A	04-11-1976
			DK 426576 A, B,	22-09-1976
			ES 405076 A	16-07-1975
			ES 405077 A	16-07-1975
			ES 405078 A	16-07-1975
			ES 405079 A	16-07-1975
			FR 2146482 A	02-03-1973
			GB 1351430 A	01-05-1974
			HU 164270 B	28-01-1974
			HU 166732 B	28-05-1975
			HU 164929 B	28-05-1974
			HU 164930 B	28-05-1974
			IE 37665 B	14-09-1977
			JP 53036471 B	03-10-1978
			LU 65753 A	28-11-1972
			NL 7210145 A, B	24-01-1973
			NL 7513437 A, B,	27-02-1976
			NO 136793 B	01-08-1977
			NO 762049 A	23-01-1973
			NO 762050 A	23-01-1973
			RO 62784 A	15-11-1977
			SE 392901 B	25-04-1977
			SE 422325 B	01-03-1982
			SE 7506819 A	13-06-1975
			US 4119635 A	10-10-1978
			ZA 7203888 A	27-06-1973
WO 9847894	A	29-10-1998	AU 6865898 A	13-11-1998
			CA 2259246 A	29-10-1998
			CN 1228089 T	08-09-1999
			EP 0922043 A	16-06-1999